

Renormalization of the Cabibbo-Kobayashi-Maskawa Matrix in Standard Model

Yong Zhou^a

^a *Institute of Theoretical Physics, Chinese Academy of Sciences, P.O. Box 2735, Beijing 100080, China*

We have investigated the present renormalization prescriptions of Cabibbo-Kobayashi-Maskawa (CKM) matrix, and found there is still not an integrated prescription to all loop levels. In this paper We propose a new prescription which is designed for all loop levels in the present perturbative theory. This new prescription will keep the unitarity of the bare CKM matrix and make the amplitude of an arbitrary physical process involving quark mixing convergent and gauge independent.

11.10.Gh, 12.15.Lk, 12.15.Hh

Since the exact examination of the Cabibbo-Kobayashi-Maskawa (CKM) quark mixing matrix [1] has been developed quickly, the renormalization of CKM matrix becomes very important [2]. This was realized for the Cabibbo angle with two fermion generations by Marciano and Sirlin [3] and for the CKM matrix of the three-generation SM by Denner and Sack [4] more than a decade ago. In recent years many people have discussed this issue [5], but a completely self-consistent prescription to all loop levels has been not obtained. In this paper we will try to solve this problem.

In general, a CKM matrix renormalization prescription should satisfy the three criterions [6]:

1. In order to keep the transition amplitude of any physical process involving quark mixing ultraviolet finite, the CKM counterterm must cancel out the ultraviolet divergence left in the loop-corrected amplitudes.
2. It must guarantee such transition amplitude gauge parameter independent [7], which is a fundamental requirement.
3. SM requires the bare CKM matrix V^0 is unitary,

$$\sum_k V_{ik}^0 V_{jk}^{0*} = \delta_{ij} \quad (1)$$

with i, j, k the generation index and δ_{ij} the unit matrix element. If we split the bare CKM matrix element into the renormalized one and its counterterm

$$V_{ij}^0 = V_{ij} + \delta V_{ij} \quad (2)$$

and keep the unitarity of the renormalized CKM matrix, the unitarity of the bare CKM matrix requires

$$\sum_k (\delta V_{ik} V_{jk}^* + V_{ik} \delta V_{jk}^* + \delta V_{ik} \delta V_{jk}^*) = 0 \quad (3)$$

Until now there are many papers discussing this problem. The modified minimal subtraction ($\overline{MS}$) prescription [8] is the simplest one, but the introduced μ^2 -dependent term is very complicated to be dealt with. In the on-shell renormalization framework, however, there is still not an integrated CKM renormalization prescription. The early prescription which used the $SU_L(2)$ symmetry of SM to relate the CKM counterterm with the fermion wave-function renormalization constants (WRC) [4,9] is a delicate and simple prescription, but unfortunately it reduces the physical amplitude involving quark mixing gauge dependent¹ [10,11]. A revised version of this prescription is to replace the on-shell fermion WRC in the CKM counterterm with the one calculated at zero momentum [10]. Another revised method is to rearrange the off-diagonal quark WRC in a manner similar to the pinch technique [11]. Different from the idea of Ref. [4], another idea is to formulate the CKM renormalization prescription with reference to the case of no mixing of fermion generations. The main idea of this prescription is to make the transition amplitude of W gauge boson decaying into up-type and down-type quarks equal to the same amplitude but in the case of no mixing of quark generations [12]. All of the mentioned prescriptions are only applied to one-loop level and are very difficult to be generalized to higher loop levels [10–12]. So we want to propose a new method to solve this problem.

¹It is easy to be understood since the $SU_L(2)$ symmetry of SM has been broken by the Higgs mechanism

We will renormalize the CKM matrix through two steps. First we introduce a CKM counterterm which makes the physical amplitude of $W^+ \rightarrow u_i \bar{d}_j$ ultraviolet convergent and gauge independent. Next we mend it to satisfy the unitary condition of Eq.(3), simultaneously keep the divergent and gauge-dependent part of it unchanged. In order to elaborate our idea clearly we firstly introduce the n-loop ($n \geq 1$) decaying amplitude of $W^+ \rightarrow u_i \bar{d}_j$ as follows (here all of the counterterms lower than n-loop level merge into the formfactors):

$$T_n = A_L[F_{Ln} + V_{ij}(\frac{\delta g_n}{g} + \frac{1}{2}\delta Z_{Wn}) + \frac{1}{2}\delta \bar{Z}_{ik}^{uLn} V_{kj} + \frac{1}{2}V_{ik}\delta Z_{kj}^{dLn} + \delta V_{ijn}] + A_R F_{Rn} + B_L G_{Ln} + B_R G_{Rn} \quad (4)$$

with g and δg the $SU(2)$ coupling constant and its counterterm, δZ_W the W boson WRC, $\delta \bar{Z}^{uL}$ and δZ^{dL} the left-handed up-type and down-type quark's WRC [13]. The added denotation "n" represents the n-loop result, and

$$\begin{aligned} A_L &= \frac{g}{\sqrt{2}} \bar{u}_i(p_1) \not{\epsilon} \gamma_L \nu_j(q - p_1), \\ B_L &= \frac{g}{\sqrt{2}} \bar{u}_i(p_1) \frac{\epsilon \cdot p_1}{M_W} \gamma_L \nu_j(q - p_1). \end{aligned} \quad (5)$$

with ϵ^μ the W boson polarization vector, γ_L and γ_R the left-handed and right-handed chiral operators, M_W the W boson mass. Similarly, replacing γ_L with γ_R in above equations we get A_R and B_R respectively. $F_{L,R}$ and $G_{L,R}$ are four formfactors. Here we only care about the coefficient of A_L which contains the n-loop CKM counterterm. The simplest method to make the amplitude T_n convergent and gauge independent is to make the coefficient of A_L equal to zero (the remaining terms should be convergent and gauge independent [10]). That's to say

$$\delta V_{ijn} = -F_{Ln} - V_{ij}(\frac{\delta g_n}{g} + \frac{1}{2}\delta Z_{Wn}) - \frac{1}{2}\delta \bar{Z}_{ik}^{uLn} V_{kj} - \frac{1}{2}V_{ik}\delta Z_{kj}^{dLn} \quad (6)$$

Obviously such CKM counterterm cannot guarantee the unitarity of the bare CKM matrix. It needs to be mended. Here we introduce a new denotation $\delta \bar{V}_n$ to denote the amended CKM counterterms which will satisfy the unitary condition of Eq.(3) to n-loop level. Our method is to by recursion construct $\delta \bar{V}_n$ through $\delta V_n, \delta \bar{V}_{n-1}, \dots, \delta \bar{V}_1$. Here we state that δV_n is obtained by using $\delta \bar{V}_{n-1}, \dots, \delta \bar{V}_1$ as the lower loop CKM counterterms in Eq.(6). Now the unitary condition of Eq.(3) becomes

$$\begin{aligned} \delta \bar{V}_1 V^\dagger + V \delta \bar{V}_1^\dagger &= 0, \\ \delta \bar{V}_2 V^\dagger + V \delta \bar{V}_2^\dagger &= -\delta \bar{V}_1 \delta \bar{V}_1^\dagger, \\ \delta \bar{V}_3 V^\dagger + V \delta \bar{V}_3^\dagger &= -\delta \bar{V}_1 \delta \bar{V}_2^\dagger - \delta \bar{V}_2 \delta \bar{V}_1^\dagger, \\ \dots\dots\dots \\ \delta \bar{V}_n V^\dagger + V \delta \bar{V}_n^\dagger &= -\delta \bar{V}_1 \delta \bar{V}_{n-1}^\dagger - \delta \bar{V}_2 \delta \bar{V}_{n-2}^\dagger \dots - \delta \bar{V}_{n-2} \delta \bar{V}_2^\dagger - \delta \bar{V}_{n-1} \delta \bar{V}_1^\dagger, \\ \dots\dots\dots \end{aligned} \quad (7)$$

In order to solve these equations, we introduce a set of symbols B_n

$$\begin{aligned} B_0 &= 0, \\ B_n &= \sum_{i=1}^{n-1} -\delta \bar{V}_i \delta \bar{V}_{n-i}^\dagger. \end{aligned} \quad (8)$$

Obviously B_n satisfies

$$B_n = B_n^\dagger \quad (9)$$

Assuming that we have obtained the counterterms $\delta \bar{V}_1, \delta \bar{V}_2, \dots, \delta \bar{V}_{n-1}$ and δV_n , the n-loop CKM counterterm $\delta \bar{V}_n$ is determined as follows:

$$\delta \bar{V}_n = \frac{1}{2}(\delta V_n - V \delta V_n^\dagger V + B_n V) \quad (10)$$

At one-loop level this result is similar as Eq.(11) of ref. [6]. It is very easy to see that such CKM counterterm satisfies Eqs.(7) to n-loop level.

The remaining problem is to test whether the amended CKM counterterm $\delta \bar{V}_n$ has the same divergent and gauge-dependent terms as δV_n , which is the requirement of making the physical amplitude involving quark mixing ultraviolet finite and gauge independent. Our answer is positive to this question. Based on the renormalizability and predictability of SM, we can predict that the divergent and gauge-dependent part of δV_n must satisfy the unitary condition of Eq.(3) at n-loop level

$$\delta V_n^{DG} V^\dagger + V \delta V_n^{DG\dagger} = B_n^{DG} \quad (11)$$

where the superscript "DG" denotes the divergent/gauge-dependent part of the quantity. This is because if not so the unitary condition of Eq.(3) will require the divergent and gauge-dependent part of the CKM counterterm different from δV_n thus will reduce the physical amplitude of $W^+ \rightarrow u_i \bar{d}_j$ divergent and gauge dependent. At one-loop level this relationship has been proven in actual calculations [6,10]. So from Eq.(10) and (11) it is obtained

$$(\delta \bar{V}_n^{DG} - \delta V_n^{DG}) V^\dagger = \frac{1}{2} (B_n^{DG} - \delta V_n^{DG} V^\dagger - V \delta V_n^{DG\dagger}) = 0 \quad (12)$$

This identity manifests that

$$\delta \bar{V}_n^{DG} = \delta V_n^{DG} \quad (13)$$

Now we have obtained the proper CKM counterterms to all loop levels, which satisfy the unitary condition of the bare CKM matrix and make the physical amplitude involving quark mixing convergent and gauge independent. We state that at present all of the CKM renormalization prescriptions are only applied to one-loop level and an integrated prescription to all loop levels doesn't appear. This situation shows that the renormalization of CKM matrix isn't an easy job. Our prescription affords a more straightforward and simple method to solve this problem and it will be easy to calculate the CKM counterterm in actual calculations (the n-loop CKM counterterm is shown in Eq.(10) and the explicit one-loop result is shown in appendix). On the other hand we suppose our prescription will not break the present symmetries of SM, e.g. Ward-Takahashi identity, because it only changes the values of CKM matrix elements from V_{ij}^0 to $V_{ij} + \delta V_{ij}$.

Acknowledgments

The author thanks professor Xiao-Yuan Li for his useful guidance and Dr. Hu qingyuan for his sincerely help (in my life).

Appendix

In this appendix we give the explicit result of $\delta \bar{V}_1$. In the on-shell renormalization framework δg and δZ_W are both real quantities, so we can obtain the following result through Eq.(10), (8) and (6)

$$\begin{aligned} \delta \bar{V}_{ij1} &= \frac{1}{2} (\delta V_{ij1} - \sum_{kl} V_{ik} \delta V_{kl1}^\dagger V_{lj}) \\ &= \frac{1}{2} (\sum_{kl} V_{ik} F_{L1lk}^* V_{lj} - F_{L1ij}) + \frac{1}{4} \sum_k (\delta \bar{Z}_{ki}^{uL1*} - \delta \bar{Z}_{ik}^{uL1}) V_{kj} + \frac{1}{4} \sum_k V_{ik} (\delta Z_{jk}^{dL1*} - \delta Z_{kj}^{dL1}) \end{aligned} \quad (14)$$

which is gauge independent since δV_{ij1} is gauge independent [10]. The ultraviolet divergence of $\delta \bar{V}_{ij1}$ is

$$\begin{aligned} \delta \bar{V}_{ij1}|_{UV} &= \frac{3\alpha\Delta}{64\pi M_W^2 s_W^2} \left[-\frac{2 \sum_{k,l \neq j} m_{d,j} m_{u,k}^2 V_{il} V_{kl}^* V_{kj}}{m_{d,l} - m_{d,j}} + \frac{2 \sum_{k,l} m_{d,j} m_{u,k}^2 V_{il} V_{kl}^* V_{kj}}{m_{d,l} + m_{d,j}} - \frac{2 \sum_{k \neq i,l} m_{u,i} m_{d,l}^2 V_{il} V_{kl}^* V_{kj}}{m_{u,k} - m_{u,i}} + \right. \\ &\quad \left. \frac{2 \sum_{k,l} m_{u,i} m_{d,l}^2 V_{il} V_{kl}^* V_{kj}}{m_{u,k} + m_{u,i}} + V_{ij} (\sum_k V_{ik} V_{ik}^* m_{d,k}^2 + \sum_k V_{kj} V_{kj}^* m_{u,k}^2 - 2m_{d,j}^2 - 2m_{u,i}^2) \right]. \end{aligned} \quad (15)$$

where α is the fine structure constant, M_W is the W boson mass, s_W is the sine of the weak mixing angle θ_W , and $\Delta = 2/(D-4) + \gamma_E - \ln(4\pi) + \ln(M_W^2/\mu^2)$ (D is the space-time dimensionality, γ_E is the Euler's constant and μ is an arbitrary mass parameter). $m_{u,i}$, $m_{d,j}$ etc. are up-type and down-type quark's masses. The R_ξ -gauge and the Dimensional regularization have been used. This result agrees with the results of Ref. [4] and [10].

- [1] N.Cabibbo, Phys. Rev. Lett. **10**, 531 (1963);
M.Kobayashi and K.Maskawa, Prog. Theor. Phys. **49** (1973) 652;
Particle Data Group, D.E. Groom et al., Eur. Phys. J. C **15** (2000) 1;
A. Höcker, H. Lacker, S. Laplace, F.Le Diberder, Eur. Phys. J. C **21** (2001) 225;
M. Beneke et al., hep-ph/0003033.
- [2] A. Sirlin, Rev. Mod. Phys. **50** (1978) 573;
W.J. Marciano and A. Sirlin, Nucl. Phys B **93** (1975) 303.

- [3] W.J. Marciano, A. Sirlin, Nucl. Phys. B**93** (1975) 303.
- [4] A. Denner, T. Sack, Nucl. Phys. B**347** (1990) 203.
- [5] J. Manzano, hep-ph/0208068;
A. Pilaftsis, Phys. Rev. D**65** (2002) 115013.
- [6] K.-P. O. Diener, B. A. Kniehl, Nucl. Phys. B**617** (2001) 291.
- [7] C. Becchi, A. Rouet, and R. Stora, Commun. Math. Phys. **42** (1975) 127; Ann. Phys. (N.Y.) **98** (1976) 287.
- [8] D.J. Gross and F. Wilczek, Phys. Rev. D**8** (1973) 3633;
W.E. Caswell and F. Wilczek, Phys. Lett. B**49** (1974) 291;
H. Kluberg-Stern and J.B. Zuber, Phys. Rev. D**12** (1975) 467;
W.A. Bardeen, A.J. Buras, D.W. Duke, T. Muta, Phys. Rev. D**18** (1978) 3998.
- [9] D. Espriu, J. Manzano, Phys. Rev D**63** 073008.
- [10] P. Gambino, P.A. Grassi, F. Madricardo, Phys. Lett. B 454 (1998) 98;
B.A. Kniehl, F. Madricardo, M. Steinhauser, Phys. Rev. D 62 (2000) 073010.
- [11] Y. Yamada, Phys. Rev. D**64** (2001) 036008.
- [12] A. Barroso, L. Brücher, R. Santos, Phys. Rev. D 62 (2000) 096003.
- [13] D. Espriu, J. Manzano, P. Talavera, Phys. Rev. D**66** (2002) 076002;
Yong Zhou, J. Phys. G, **29** (2003) 1.